



Mole Cricket Bait

A CRUMBILIZED BAIT AND CONTACT INSECTICIDE FOR CONTROL OF FIELD CRICKETS, MOLE CRICKETS, AND MANY OTHER INSECTS AS LISTED INFESTING GOLF COURSE TURF, WAREHOUSES, FOOD PROCESSING PLANT SITES, INDUSTRIAL PLANT SITES, AND MANUFACTURING PLANT SITES.

CONTAINS AN ATTRACTANT

ACTIVE INGREDIENTS:

Chlorpyrifos (0,0-diethyl 0-(3, 6, 6-trichloro-2-pyridyl) phosphorothioate).....	1.0%
Other Ingredients:	99.0%
TOTAL	100.0%

50 POUNDS NET WEIGHT

EPA Reg No. 769-679

EPA Est. No. 44616-GA-1 (C)

EPA Est. No. 44616-MO-1 (C)

Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

If Swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. DO NOT induce vomiting or give anything by mouth to an unconscious or convulsing person. Get medical attention.

If In Eyes: Flush eyes with plenty of water.

If on Skin: In case of contact, remove contaminated clothing and wash affected area with soap and water.

Note To Physician: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote only by injection.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed. Avoid contact with eyes, skin and clothing. Avoid breathing dust. Wash thoroughly after handling and before eating and smoking. Keep away from food, feedstuffs, and water supplies.

PERSONAL PROTECTIVE EQUIPMENT(PPE)

Applicators and Other Handlers Must Wear: Long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks and a Dust/Mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N, R, P, or HE filter.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water, to areas where surface is present or to intertidal areas below the mean high water mark. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. Use only as directed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store product in its original labeled container in a cool, dry, locked place out of reach of children.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

IMPORTANT: Read these entire Directions and Conditions of Sale before using this product. **DISCLAIMER:** Value Garden Supply warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purpose referred to in the Directions For Use, subject to the inherent risks referred to above. Value Garden Supply makes no implied warranty. In no case shall Value Garden Supply be liable for consequential, special, or indirect damages resulting from the use, handling or application of this product.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Mole Crickets In Golf Course Turf: Apply this product uniformly using any hand or power operated gravity or rotary type spreader designed for application of granules at the rate of 2-1/4 lbs. per 1,000 sq. ft. (equivalent to 100 lbs. per acre). For best results if insect populations are serious, plan on a second treatment in about 7 to 10 days.

Golf Course Turf – Surface Feeding Insects: Use 2-1/4 (1,000 sq. ft.) for control of Ants, Armyworms, (such as Beet, Fall, Yellowstriped), Centipedes, Chinch bugs, Cutworms, Crickets, Earwigs, Grasshoppers, Millipedes, Sod Webworms (lawn moth), and Sowbugs. For Sod webworms also apply at rate of 2-1/4 per 1,000 sq. ft., but delay watering or mowing the treated area for 12 to 24 hours after treatment. Treat in the spring or when insects first appear troublesome. Additional treatments at 30 day intervals may be needed for new infestations. It is not necessary to water the insecticide in to control these surface insects.

Crawling Insect Control Around Warehouses, Food Processing Plant Sites, Industrial Plant Sites, and Manufacturing Plant Sites: To protect against insects entering the indoors, a protective band of this product granules may be placed around the outside of buildings to control Ants, Cockroaches, Field and Mole Crickets, Earwigs, Millipedes, Sowbugs and Springtails before they attempt to enter. Use 1 pound to treat 440 sq. ft. (as a band 5 ft. x 88 ft. long) around the structure. Apply when insects have become troublesome or before they cause problems in the spring or early fall. Repeat one month later and then as needed for control, but not more than once every 7 days.

For Professional Use Only. Imported Fire Ants – Individual Mound

Treatment: For public health treatment of Fire Ant mounds occurring in lawns, turf and premise areas, treat individual ant mounds measuring from 3 to 12 inches in diameter by applying 8 tablespoonfuls (1.6 oz.) of granules evenly in a circle around the outer edges (1 to 2 feet from the center) of the mound. Finish the treatment by applying 5 tablespoonfuls (1 oz.) of sprinkled evenly over the center of the mound. Best results will be obtained if mound is not disturbed after treatment thus allowing ants to come and go and to be exposed to the insecticide. Repeat treatments as needed for additional control, but not more than once every 7 days.

SPREADER SETTINGS

SPREADER	2 1/4 LBS. OF PRODUCT (1,000 SQ. FT.)
Cyclone B 1	#4-1/2



Distributed by:
Value Garden Supply, LLC
PO Box 585
St. Joseph, MO 64502-0585
(952) 884-6477
www.valuegardens.com

RM22010