



Indoor/Outdoor Bug Killer Spray

For Control of Pests on Livestock.

For Control Of Fleas, Brown Dog Ticks and Lice On Premises.

Provides 60-Day Control of Cockroaches, Fleas and Stored Product Pests.

For Domestic Use Both Indoors and Outdoors.

ACTIVE INGREDIENTS:

*Permethrin 0.25%

INERT INGREDIENTS: 99.75%

TOTAL 100.0%

*(3-phenoxyphenyl) methyl (+/-) cis/trans 3-(2,2-dichloroethenyl) 2,2-dimethylcyclopropanecarboxylate.

Cis/trans ratio: min.35%(+)cis and max. 65% (+/-) trans.

EPA Registration No. 572-350

EPA Establishment No. 44616-MO-1

KEEP OUT OF REACH OF CHILDREN CAUTION

READ ENTIRE LABEL BEFORE EACH USE

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Wash thoroughly after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before surface applications. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

ENVIRONMENTAL HAZARDS: This product is extremely toxic to fish. Do not apply directly to any body of water. Do not contaminate water when disposing of equipment washwaters.

FIRST AID

If Swallowed	<ul style="list-style-type: none"> Have person sip a glass of water if able to swallow. Call a poison control center or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If on Skin or Clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If In Eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

For information regarding medical emergencies or pesticide incidents, call the International Poison Center at 1-888-740-8712

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area above 32°.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

CONTAINER DISPOSAL: If Empty, do not reuse this container. Place in trash or offer for recycling if available.

If Partly filled, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not use on plants being grown for sale or other commercial seed production or research purposes.

Do Not Use on Dogs or Cats.

SHAKE WELL BEFORE USING.

INDOOR APPLICATION

For spot application only. Do not use as a space spray indoors. In the home, cover all food handling surfaces and cover or remove all food and cooking utensils, or wash thoroughly after treatment. Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

Surface Spraying : (Ready To Use) For 60 day (2 months) control of Cockroaches, Waterbugs, Palmetto Bugs, Scorpions, Millipedes, Centipedes, Sowbugs, Pillbugs, Ants, Silverfish, Firebrats, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Drugstore Beetles, Meal Worms, Grain Mites and Cadelles: Direct the spray into hiding places, cracks and crevices and behind shelves and drawers. For Silverfish, spray bookcases. For Ants, spray trails, nests and points of entry.

To Control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, crevices and infested areas of shelving.

To Control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors.

To Control Fleas (Adult and Larvae) and Ticks (Adult and Larvae): Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind

baseboards, moldings, window and door frames, and entire areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Allow to dry thoroughly before allowing pets to contact treated areas. Do Not Use on Cats or Dogs.

To Control Clothes Moths: Remove any infested articles from storage, brush thoroughly, and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints, and crevices. After airing, infested articles may be treated lightly, keeping nozzle at least three feet from fabric to avoid staining.

To Kill Lice and Louse Eggs: Spray in an inconspicuous area to test for possible staining or discoloration. Inspect again after drying, then proceed to spray entire area to be treated. Spray from a distance of 8 to 10 inches. Treat only those garments and parts of bedding, including mattresses and furniture that cannot be either laundered or dry cleaned. Allow all treated articles to dry thoroughly before use.

LIVESTOCK PESTS

Ready to use. Affix trigger, pump or other type sprayer to container for application. Do not transfer to another container.

ON LIVESTOCK: To protect cattle (beef and dairy), goats, sheep, hogs and horses from Horn Flies, Face Flies, House Flies, Mosquitoes and Gnats: Apply a light mist sufficient to set the surface of the hair. For face flies, spray face and head, but do not spray into eyes. **To Control Stable Flies, Horse Flies and Deer Flies:** Apply at a rate of 2 ounces per adult animal, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection. **To Control Blood-Sucking Lice:** Apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 3 weeks if required. **To Control Poultry Lice:** Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist. **To Control Bedbugs and Mites in Poultry Houses:** Spray roosts, walls and nests or cages thoroughly. This should be followed by spraying over the birds with a fine mist. Spray crevices of roost poles, cracks in walls, and cracks in nests where the bedbugs and mites hide.

OUTDOOR APPLICATION

To help control ticks in outdoor areas: Ready to use. Treat a band of vegetation and leaf litter 6-10 feet wide around the outside of buildings and adjacent to activity areas, especially where dense vegetation occurs.

Pests On The Outdoor Surfaces of Buildings: Dilute before using. To temporarily control House Flies, Mosquitoes, Cluster Flies, Gnats and Flying Moths. For outdoor use only as an aid in reducing annoyance from these insects. Dilute before using. Use clean sprayer. Carefully measure and mix 1 fluid ounce of AllPro Indoor/Outdoor Bug Killer Spray and 12 fluid ounces of water. Using dilute mixture, spray pests on outside surfaces of screens, doors, window frames or wherever these insects may enter the room. Also treat surfaces around light fixtures on porches, in garages and other places where these insects alight or congregate. Flush sprayer with clean water after each use.

ORNAMENTAL FLOWER GARDENS: Ready to use. Affix trigger, pump or other type of sprayer to container for application. Do not transfer to another container.

Garden Insects: For use on roses, dahlias, asters and other ornamentals to kill Japanese Beetles hit by spray as well as Whiteflies, Aphids, Exposed Thrips, Spiders, Red Mites and Leafminers. May be used on plants as a multi-purpose spray in small gardens and for spot treatments of incipient insect infestations to prevent spreading in large garden areas. May also be used on other plants such as African violets, asters, azaleas, begonias, camellias, carnations, chrysanthemums, dahlias, delphiniums, dogwoods, English ivy, eunonymus, fuschia, geraniums, crassula, grape vines, Kentia palms, laurels, marigolds, rhododendrons, roses, rubber plants, snapdragons, stocks, wandering jews and zinnias.

Effective when used as directed against Whiteflies, Aphids, Spider and Red Mites, exposed Thrips, Lace Bugs, Japanese Beetles, Pavement Ants, Clover Mites and Armyworms. When spraying plants, do not operate closer than 18 inches. Use sweeping motion. Be cautious about wetting tender foliage, young plants and new growth. Do not spray plants when temperatures exceed 90 F. Contact insects directly whenever possible. Repeat treatment as infestations occur.

WARRANTY: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 572-350 EPA Est. No. 44616-MO-1

Value Garden Supply

P.O. Box 585

St. Joseph, MO 64502-0585

(952) 884-6477

www.valuegardens.com