



**Gardener's  
Choice™**

Ready To Use

**Kills Grubs, Ants,  
Fleas, Brown Dog  
Ticks, Spiders,  
and Other  
Insects Listed**

**Provides Contact  
and Residual Control  
For Up To 4 Weeks**


**NET CONTENTS  
10 lbs. (4.53 kg)**



**Gardener's  
Choice™**

# LAWN INSECT

## Killer



**ACTIVE INGREDIENT:**

\*Permethrin ..... 0.25%

INERT INGREDIENTS: ..... 99.75%

TOTAL ..... 100.00%

\*Cis/trans ratio: Maximum 55% (±) cis and minimum 45% (±) trans.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION** See Additional Precautionary  
Statements on back of bag.

# LAWN INSECT Killer



# Gardener's Choice™ Lawn Insect Killer

## SPOT TREATMENT

FIRST AID	
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

### PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling. Avoid contamination of food or feed. Utensils such as teaspoons and measuring cups should not be used for food purposes after use with insecticides.

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not apply directly to water. Drifts and runoffs from treated areas may be hazardous to aquatic organisms in treated areas. Do not contaminate water when disposing of equipment washwaters.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with pesticides.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

**Gardener's Choice Lawn Insect Killer** is recommended for the control of lawn insects, fleas, and ticks outdoors in residential areas. Repeated applications are made by broadcast treatment with a spreader to lawns, under shrubbery, and other ornamental and turf grass areas, band treatments are recommended around house foundations and spot applications are made to ant mounds and wasp nests.

### THREE EASY STEPS TO USE THE PRODUCT

- Step #1 Determine size of area to be treated - measure and multiply length times width.
  - Step #2 Determine appropriate spreader setting using chart
  - Step #3 Treat as directed - thorough coverage is important. Water in where indicated.
- Keep children and pets off treated areas until this material is washed into turf and grass is dry.

### OVERALL LAWN TREATMENT FOR GRASS AND DICHONDRA LAWNS

**PESTS:** Ants, Armyworms (inc. Fall, Yellow-striped, ect.), Bermudagrass mites, Boxelder Bugs, Centipedes, Chiggers, Chinch Bugs, Clover Mites, Cockroaches, Crickets, Cutworms, Darkling Ground Beetles, Digger wasps, Essex Skipper, European Crane Fly Larvae, Field Crickets, Fiery Skipper, Flea Beetle (In dichondra), Grasshopper, Hyperodes (Turfgrass Weevil), Lawn Moths, Leafhoppers, Scorpions, Silverfish & Firebrats, Sod Webworms, Sowbugs, Spiders (Inc. Black Widow), Spittlebugs, Springtails, Vegetable Weevils (In dichondra).

**USE RATE:** Treat when insects first appear using granules at a rate of 2 pounds per 1,000 sq. ft. or, for heavy lawns such as St Augustine grass, 3 pounds per 1,000 sq. ft. using a fertilizer spreader. See spreader setting chart below.

**HOW TO USE:** Apply this product uniformly to home lawns and under shrubbery. To help obtain uniform distribution, apply one-half required amount of the product in one direction, then apply the balance at a right angle. Immediately after application, water the lawn lightly. European Crane fly larvae (Leatherjackets) - the best time to apply is in May or June when the insects are young. Chinch bugs - Start treatments when temperatures reach 80° F or above. A second application may be necessary in 6 weeks if insect activity continues.

**PESTS:** Billbugs, Grubs, Japanese Beetle (adult), Dung Beetles (Ateenius Spretulus)

**USE RATE:** Use granules at the rate of 2 1/2 pounds per 1,000 sq. ft.

**HOW TO USE:** Water grass thoroughly after application. For Billbug control, apply when activity is first observed or when chewed or brown grass indicates damage from these insects. For grub control, except Dung Beetle, treat any time between late July and early October. For Dung Beetle Grub control, apply in spring or early summer when damage is first noticed. Water grass thoroughly with 1/4 to 1/2 inch of water after application. Treat again in late summer or early fall if a second generation occurs.

### BAND TREATMENT AROUND HOUSE FOUNDATION

**PESTS:** Ants, Boxelder Bugs, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Field Crickets, Fleas, Scorpions, Sowbugs, Spiders (including Black Widow), Springtails (Collembola) and Ticks.

**USE RATE:** 1 lb. per 500 sq. ft. or 2 lbs. per 1,000 sq. ft.

**HOW TO USE:** To help prevent infestation of buildings, apply a 5 foot band of granules next to the house foundation. Distribute granules uniformly. Water lightly after application.

Distributed by:  
Value Garden Supply  
PO Box 585  
St. Joseph, MO 64502-0585  
(952)884-6477

EPA Reg. No. 28293-328-192  
EPA Est. No. 44616-MO-1(14) 44616-GA-1(14)  
Superscript used is first letter of lot number.

RM#17079

### ANTS NESTING OUTDOORS INCLUDING

Acrobat, Argentine, Big Headed, Cornfield, Field, Harvester, Leafcutter, Little Black, Pavement and Pyramid Ants.

**USE RATE:** Product should be applied to thoroughly cover the nest (1 to 2 teaspoons) and sprinkled gently with water.

**HOW TO USE:** If possible, it is best to try to locate nest and treat this first. Nests will usually be found in the ground but may be concealed under stones, boards, slabs or in stumps or logs. In addition to treating ants nests or mounds, entry points of buildings, (used by ants) and the soil surface along the perimeter of the building should be treated.

### DIGGER WASPS:

**USE RATE:** Apply 1 to 2 teaspoons of granules in and around each individual ground nest opening.

**HOW TO USE:** Make application at dusk when the wasps are not active.

### IMPORTED FIRE ANTS:

**USE RATE:** Apply one cup over, and 2 feet around each fire ant mound.

**HOW TO USE:** Immediately after application, water the granules into the soil gently with a light sprinkling to avoid disturbing the colony. Use at least 1 gallon of water per mound, but no more than 2 gallons. Use of high pressure watering equipment may disturb the colony and cause migration of the ants, thus reducing the effectiveness of the treatment. Apply in cool weather (65 to 80° F), or in early morning or late evening for best results. Treat new mounds as they appear.

### FLEA AND TICK CONTROL (OUTDOORS)

In summer months, fleas and ticks frequently become established in lawns and pet resting areas surrounding the home. For best control of Fleas and Ticks (including but not limited to Bear Tick, Black-legged Tick, Brown Dog Tick, Deer Tick (Ixodes dammini), Lone Star Tick, Western Black-legged Tick, Wood Tick): a regular treatment program should be followed. Treat once a month when pests first appear in late spring (May-June) and continue until late summer (September-October). Fall temperatures below 50° F decrease pest activity. Eliminating weeds around the home and mowing lawns frequently reduces the hiding places for these pests.

Note: Dogs, cats their kennel or bedding area should also be treated with a registered product. In addition, if fleas and ticks are indoors on carpets or pet resting areas, use a registered product ideally in combination with an Insect Growth Regulator. It is very important that all three sites - outdoors, indoors and also the pet be treated with appropriate labeled products to achieve satisfactory control. If any of these areas is not treated, it could serve as a source of reinfestation to the entire premises.

### SOD WEBWORMS (CRAMBUS SPP)

Sod Webworms are the light brown caterpillars 1/4 to 3/4 inch long, that hide in the grass during the day and feed on the grass blades at night. Adult Webworms, usually referred to as Lawn Moths (about 1 inch in length with long snout) emerge from the grass at dusk. They fly over the lawn laying eggs that later hatch into destructive caterpillars. Damage appears first as small brown spots which, if not treated, may rapidly develop into large brown areas of severely damaged or dead grass. If Lawn Moths (Sod Webworm Adults) are noticed in large numbers at dusk, or at first sign of damage to lawn, treat entire lawn evenly.

### SPREADER SETTINGS

Spreader Make and Model	Grass or Dichondra Lawns at 2 lbs. per 1000 sq. ft.	White Grubs, Billbugs at 2 1/2 lbs. Per 1000 sq. ft.	St. Augustine Lawns at 3 lbs. per 1000 sq. ft.
Cyclone Broadcast	3	4	5
Greenview Drop Spread III	3	4	5
Greenview Jet Spred III	4	5	6
Republic Drop EZ 530	6 1/2	7	8
Republic Broadcast EZ 540	2	3	4
Scotts AG3	6	6 1/2	7
Scotts SG3	4	4 1/2	5
Scotts EG3	24	25	26
Scotts PF4	6	6 1/2	7
Sears Drop 1927	5 1/2	6	7
Sears Broadcast 1922	5 1/2	6	7

Note: Spreader settings given are approximate and variation can occur between identical models. User should calibrate spreader by accurately weighing a few pounds into your spreader and applying to a measured area. Then adjust setting if necessary to assure correct rate of application.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Information Center at 1-800-858-7378.

### STORAGE AND DISPOSAL

**Storage:** Store product in its original container in a cool, dry locked place out of reach of children.  
**Disposal: If Empty** - Do not reuse this container. Place in trash or offer for recycling if available.  
**If Partly Filled** - Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

### LIMITED WARRANTY AND DISCLAIMER

Buyer assumes all responsibility for safety and use not in accordance with directions.

